

Zidovudine

Brand Name: Retrovir



Drug Description

Zidovudine is a synthetic pyrimidine nucleoside analogue of the naturally occurring nucleoside thymidine. [1]

HIV/AIDS-Related Uses

Zidovudine was approved by the FDA on March 19, 1987, for the treatment of HIV infection in combination with other antiretroviral agents. Studies indicate that zidovudine in combination with other antiretroviral agents is superior to monotherapy for one or more of the following endpoints: delaying death, delaying development of AIDS, increasing CD4 cell counts, and decreasing plasma HIV RNA.[2] [3]

Because monotherapy with any single antiretroviral agent is no longer considered an acceptable option in the treatment of HIV infection, except in the prevention of perinatal transmission of HIV from an infected mother to her child, zidovudine is used in conjunction with other antiretroviral agents for initial antiretroviral therapy in the management of HIV infection in treatment-naïve patients. When zidovudine is included in an antiretroviral regimen that includes other nucleoside reverse transcriptase inhibitors (NRTIs), didanosine or lamivudine is usually preferred and zalcitabine is considered an alternative. Although two-drug regimens that include only NRTIs are no longer preferred regimens for previously treated adults, several studies have evaluated the efficacy of other two-drug and three-drug regimens containing zidovudine in treatment-experienced patients. Results indicate that regimens that include zidovudine and lamivudine or an HIV protease inhibitor (e.g., indinavir, ritonavir, saquinavir) generally are more effective than zidovudine monotherapy in increasing CD4 counts and decreasing plasma HIV-1 RNA levels in treatment-experienced patients. Zidovudine has also been used in conjunction with nevirapine, a nonnucleoside reverse transcriptase inhibitor (NNRTI), as part of treatment for previously treated HIV infected adults.[4]

Zidovudine is also approved for the prevention of

vertical transmission of the HIV virus from an HIV-infected mother to her fetus, as part of a regimen that includes oral zidovudine administered to the mother beginning at 14 to 34 weeks of gestation, intravenous zidovudine administered to the mother during labor, and zidovudine administered to the neonate for the first 6 weeks of life. However, transmission to infants may still occur in some cases despite the use of this regimen.[5]

Zidovudine is used in conjunction with lamivudine for postexposure prophylaxis of HIV infection in healthcare workers and other individuals exposed occupationally via percutaneous injury or mucous membrane or nonintact skin contact with blood, tissues, or other body fluids associated with a risk for HIV transmission.[6]

Pharmacology

Zidovudine is virustatic, acting as a reverse transcriptase inhibitor. Zidovudine is phosphorylated intracellularly to its active 5'-triphosphate metabolite, zidovudine triphosphate (ZDV-TP), by cellular kinases. Neither zidovudine itself nor its intermediate monophosphate metabolite has in vitro activity against HIV. Further study is needed to determine if the intermediate diphosphate metabolite has antiretroviral activity. Because phosphorylation of zidovudine depends on cellular enzymes rather than viral enzymes, conversion to the active triphosphate derivative occurs in HIV infected and uninfected cells. Following conversion, the pharmacologically active metabolite inhibits in vitro replication of HIV by interfering with the viral RNA-directed DNA polymerase, reverse transcriptase (RT). ZDV-TP appears to compete with thymidine triphosphate for incorporation into viral DNA by the RT enzyme. After incorporation of ZDV-TP, DNA synthesis is prematurely terminated because the 3'-azido group in the zidovudine molecule prevents further 5' to 3' phosphodiester linkages.[7]

Zidovudine is absorbed rapidly and almost completely from the gastrointestinal tract, with peak serum concentrations occurring in adults within 0.4 to 1.5 hours after an oral dose.

Zidovudine

Pharmacology (cont.)

Zidovudine appears to undergo first-pass metabolism. In fasting adults, about 65% of an oral dose reaches systemic circulation as unchanged drug. The peak serum concentration achieved following administration of zidovudine tablets is equivalent to that following administration of capsules or oral solution; however, absorption following oral administration shows considerable individual variability.[8]

There is limited information on the distribution of zidovudine in the body, but the drug appears to be widely distributed. The apparent volume of distribution for the drug in adults with HIV infection is 1.4 to 1.6 L/kg. Zidovudine is distributed into the cerebrospinal fluid (CSF) following both oral or intravenous (IV) administration. The ratio of CSF/plasma concentration of zidovudine reported in various studies in adults or children with HIV infection who have received oral or IV therapy with zidovudine has ranged from 0.15 to 2.1.[9] Time to peak concentration in serum is 0.5 to 1.5 hours; time to peak concentration in CSF is 1 hour after the end of a 1-hour IV infusion.[10]

Zidovudine is in FDA Pregnancy Category C. There are no adequate and well-controlled studies conducted on pregnant women. In rats, 3000 mg/kg per day (resulting in peak plasma concentrations of 350 times the peak human plasma concentration) caused marked maternal toxicity and an increase in the incidence of fetal malformations. Teratogenic effects were not seen in this experiment at doses of 600 mg/kg per day or less.[11] Zidovudine crosses the placenta with a cord-to-maternal blood ratio of approximately 0.8. Zidovudine and its glucuronide metabolite cross the human placenta and are distributed into cord blood and amniotic fluid as well as fetal liver and muscle. There is minimal distribution into fetal central nervous system (CNS) tissue. The drug is also distributed into human milk. In a study with HIV infected women who received a single 200-mg dose of zidovudine, the concentration of the drug in milk were similar to concurrent serum concentrations.[12] To monitor maternal-fetal outcomes of pregnant women exposed to zidovudine (and/or other antiretrovirals), an Antiretroviral Pregnancy

Registry has been established. Physicians may register patients at 1-800-258-4263.[13]

Plasma protein binding of zidovudine is low (30% to 38%).[14] Zidovudine is rapidly metabolized via glucuronidation in the liver principally to

Zidovudine



Pharmacology (cont.)

Following hepatic metabolism, elimination of zidovudine is primarily renal. In adults, approximately 14% to 18% is secreted by glomerular filtration and active tubular secretion into urine. Approximately 60% to 74% of the GZDV metabolite is recovered in urine within 6 hours of administration. The plasma half-life of zidovudine in adults averages approximately 0.5 to 3 hours following oral or IV administration. Following IV administration, plasma concentrations decline in a biphasic manner; half-life in adults is less than 10 minutes in the initial phase and one hour in the terminal phase. Current data on the efficacy of dialysis on removing zidovudine vary, but hemodialysis and peritoneal dialysis appear to have a negligible effect. Hemodialysis does enhance the elimination of GZDV; however, dialysis clearance of GZDV is minimal compared to the clearance of GZDV in patients with normal renal function.[16] [17]

Emergence of zidovudine resistance appears to be a function of the duration of zidovudine therapy, the severity of HIV disease, and the overall potency of the regimen in which the drug is used. Resistance is most likely to develop in patients with advanced HIV infection, those with low initial absolute helper/inducer T cell counts, and those receiving prolonged zidovudine therapy. Although it has been suggested that zidovudine resistance may develop at a slower rate in patients with asymptomatic HIV infection than in those with more advanced disease, high-level zidovudine resistance has emerged in patients with asymptomatic infection, especially in those who have received up to 3 years of zidovudine monotherapy.[18]

Although the mechanisms of resistance or reduced susceptibility to NRTIs have not been fully determined to date, specific mutations of HIV RNA-directed DNA polymerase (RT) at critical codons on the pol gene fragment have been associated with zidovudine resistance. Zidovudine resistance develops in a progressive, stepwise manner, and each reduction in susceptibility appears to be associated with the acquisition of an additional mutation in the HIV RT gene. The degree of resistance appears to depend on the number and combinations of these mutations.[19]

Further study is needed to more fully evaluate the

Zidovudine



Pharmacology (cont.)

extent of cross-resistance among the NRTIs. Some in vitro studies indicate that zidovudine-resistant HIV generally is susceptible to didanosine, zalcitabine, and stavudine; however, some zidovudine-resistant strains may also be cross-resistant to or have decreased susceptibility to other NRTIs, including didanosine, lamivudine, stavudine, and/or zalcitabine. The mutation at position 151 appears to play an important role in the development of multidrug resistance. The pattern of mutations with combination therapy is different from that seen with zidovudine monotherapy. Because the drugs have different target enzymes, cross-resistance between zidovudine and HIV protease inhibitors is unlikely. The potential for cross-resistance between zidovudine and NNRTIs is also considered to be low because the drugs bind at different sites on the RT enzyme and have different mechanisms of action.[20]

Adverse Events/Toxicity

Zidovudine has been associated with hematologic toxicity, including neutropenia, leukopenia, and severe anemia, particularly in patients with advanced HIV disease. Prolonged use of zidovudine has been associated with symptomatic myopathy. Lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported with the use of nucleoside analogues alone or in combination, including zidovudine and other antiretrovirals.[21]

The most frequent adverse effects of zidovudine are granulocytopenia and anemia. These are inversely related to the CD4 lymphocyte count at the start of therapy and directly related to dosage and duration of therapy. Significant anemia most commonly occurs after 4 to 6 weeks of therapy. Other adverse effects include changes in platelet count, hepatotoxicity, lactic acidosis, myopathy, neurotoxicity, severe headache, insomnia, myalgia, nausea, changes in pigmentation, hyperpigmentation of nails, and bone marrow depression.[22]

Redistribution/accumulation of body fat, including central obesity, dorsocervical fat enlargement

(buffalo hump), peripheral wasting, facial wasting, breast enlargement, and "cushingoid appearance," have been observed in patients receiving antiretroviral therapy. The mechanism and long-term consequences of these events are currently unknown. A causal relationship has not been established.[23]

In monotherapy clinical studies using zidovudine, the most common adverse events reported were headache, malaise, anorexia, and nausea.[24]

Drug and Food Interactions

Concurrent use of blood dyscrasia-causing medications, other bone marrow depressants, and radiation therapy with zidovudine may cause an additive or synergistic myelosuppression requiring dosage reduction of either or both drugs. Concurrent use of clarithromycin and zidovudine results in lower peak serum concentrations and delayed time to peak concentration of zidovudine.[25]

Concurrent administration of ganciclovir or interferon alfa with zidovudine is not recommended because severe hematologic toxicity may occur. Concurrent administration of these medications should be monitored frequently for abnormalities in hemoglobin, hematocrit, and white blood cell count; dose reduction or discontinuation of one or both of the medications may be necessary.[26]

Concurrent use of probenecid with zidovudine increases serum concentrations and prolongs elimination half-life for zidovudine, resulting in increased risk of toxicity. In one small trial, a very high incidence of rash was observed in patients receiving probenecid concurrently with zidovudine. Influenza-like symptoms, such as myalgia, malaise, and fever, have also occurred.[27]

Concurrent use of doxorubicin or ribavirin and zidovudine is not recommended; in vitro studies indicate an antagonistic relationship between doxorubicin or ribavirin with zidovudine. Ribavirin inhibits the phosphorylation of zidovudine to its active triphosphate form, thus antagonizing the in vitro antiviral activity of zidovudine against HIV. These drugs should not be used concurrently.[28] [29]

Zidovudine



Drug and Food Interactions (cont.)

Phenytoin plasma levels have been reported to be low in some patients receiving zidovudine. A pharmacokinetic interaction study showed no effect on phenytoin kinetics, but a 30% decrease of zidovudine clearance was observed with concurrent use of phenytoin and zidovudine.[30] [31]

Total serum concentrations of zidovudine increase when atovaquone, fluconazole, methadone, probenecid, and valproic acid are coadministered with zidovudine. Nelfinavir, rifampin, and ritonavir coadministered with zidovudine decreases the total serum concentration of zidovudine.[32]

Stavudine should not be used concurrently with zidovudine because an antagonistic relationship has been demonstrated in vitro.[33]

Zidovudine should not be administered concurrently with combination products (Combivir, Trizivir) that also contain zidovudine.[34]

Contraindications

Zidovudine is contraindicated in patients who have potentially life threatening allergic reactions to any of the components of the formulations. Zidovudine should not be administered concomitantly with any combination product tablets that contain zidovudine as one of their components.[35]

Clinical Trials

For information on clinical trials that involve Zidovudine, visit the ClinicalTrials.gov web site at <http://www.clinicaltrials.gov>. In the Search box, enter: Zidovudine AND HIV Infections.

Dosing Information

Mode of Delivery: Oral (capsules, oral solution, and tablets); intravenous injection.[36]

Dosage Form: Capsules containing 100 mg zidovudine; oral solution containing 50 mg zidovudine /5 ml in 240 ml bottle; film-coated tablets containing 300 mg zidovudine.[37] [38]

IV infusion containing 10 mg/ml in 20-ml

single-use vials.[39]

The recommended oral dose of zidovudine is 600 mg daily (either 300 mg twice daily or 200 mg three times a day). The recommended dose in pediatric patients age 6 weeks to 12 years is 160 mg/m² every 8 hours (480 mg/m²/day up to a maximum of 200 mg every 8 hours). The recommended dosing regimen for administration to pregnant women (more than 14 weeks of pregnancy) is 100 mg five times per day until the start of labor. During labor and delivery, IV zidovudine should be administered at 2 mg/kg (total body weight) over 1 hour followed by a continuous IV infusion of 1 mg/kg/hour (total body weight) until clamping of the umbilical cord. The neonate should receive 2 mg/kg orally every 6 hours starting within 12 hours after birth and continuing through 6 weeks of age. In patients maintained on hemodialysis or peritoneal dialysis, the recommended dose of zidovudine is 100 mg every 6 to 8 hours.[40]

Storage: Zidovudine capsules and tablets should be stored at 15 C to 25 C (59 F to 77 F) and protected from light. Capsules may become discolored or brittle as a result of heat and sunlight exposure, so capsules should be protected from light, heat, and moisture.[41] [42]

Zidovudine oral solution should be stored at 15 C to 25 C (59 F to 77 F).[43] [44]

Zidovudine for injection concentrate for IV infusion should be stored at 15 C to 25 C (59 F to 77 F) and protected from light.[45] [46]

Chemistry

CAS Name: Thymidine, 3'-azido-3'-deoxy-[47]

CAS Number: 30516-87-1[48]

Molecular formula: C₁₀H₁₃N₅O₄[49]

C44.94%, H4.90%, N26.21%, O23.95%[50]

Molecular weight: 267.24[51]

Melting point: 106-112 C[52]

Zidovudine



Chemistry (cont.)

Physical Description: White to beige, odorless, crystalline solid.[53]

Stability: After dilution, IV solutions are physically and chemically stable for 24 hours at room temperature, 15 C to 25 C (59 F to 77 F), and for 48 hours if refrigerated at 2 C to 8 C (36 F to 46 F). However, due to the risk of microbial contamination, diluted solutions should be administered within 8 hours if stored at 25 C (77 F) or within 24 hours if refrigerated at 2 C to 8 C (36 F to 46 F).[54]

Solubility: 20.1 mg/ml in water at 25 C.[55]

Other Names

Azidothymidine[56]

AZT[57]

3'-Azido-3'-deoxythymidine[58]

ZDV[59]

Zidovudina[60]

Further Reading

Bauer GR, Welles SL, Colgrove RR, Pitt J; Women and Infants Transmission Study Team. Zidovudine resistance phenotype and risk of perinatal HIV-1 transmission in zidovudine monotherapy-treated mothers with moderately advanced disease. *J Acquir Immune Defic Syndr*. 2003 Nov 1;34(3):312-9. PMID: 14600578

McComsey G, Maa JF. Host factors may be more important than choice of antiretrovirals in the development of lipoatrophy. *AIDS Read*. 2003 Nov;13(11):539-42, 559. PMID: 14649623

Robbins GK, De Gruttola V, Shafer RW, Smeaton LM, Snyder SW, Pettinelli C, Dube MP, Fischl MA, Pollard RB, Delapenha R, Gedeon L, van der Horst C, Murphy RL, Becker MI, D'Aquila RT, Vella S, Merigan TC, Hirsch MS; AIDS Clinical Trials Group 384 Team. Comparison of Sequential Three-Drug Regimens as Initial Therapy for HIV-1

Infection. *N Engl J Med*. 2003 Dec 11;349(24):2293-2303. PMID: 14668455

Shafer RW, Smeaton LM, Robbins GK, De Gruttola V, Snyder SW, D'Aquila RT, Johnson VA, Morse GD, Nokta MA, Martinez AI, Gripshover BM, Kaul P, Haubrich R, Swingle M, McCarty SD, Vella S, Hirsch MS, Merigan TC; AIDS Clinical Trials Group 384 Team. Comparison of Four-Drug Regimens and Pairs of Sequential Three-Drug Regimens as Initial Therapy for HIV-1 Infection. *N Engl J Med*. 2003 Dec 11;349(24):2304-2315. PMID: 14668456

Manufacturer Information

Zidovudine
GlaxoSmithKline
5 Moore Drive
Research Triangle Park, NC 27709
(888) 825-5249

Retrovir
GlaxoSmithKline
5 Moore Drive
Research Triangle Park, NC 27709
(888) 825-5249

For More Information

Contact your doctor or an AIDSinfo Health Information Specialist:

- Via Phone: 1-800-448-0440 Monday - Friday, 12:00 p.m. (Noon) - 5:00 p.m. ET
- Via Live Help: http://aidsinfo.nih.gov/live_help Monday - Friday, 12:00 p.m. (Noon) - 4:00 p.m. ET

Zidovudine



References

1. GlaxoSmithKline - Retrovir Prescribing Information, April 2003, p. 2. Available at: http://us.gsk.com/products/assets/us_retrovir.pdf. Accessed 06/15/04.
2. FDA - Drugs Used in the Treatment of HIV Infection. Available at: <http://www.fda.gov/oashi/aids/virals.html>. Accessed 06/15/04.
3. GlaxoSmithKline - Retrovir Prescribing Information, April 2003, p. 9. Available at: http://us.gsk.com/products/assets/us_retrovir.pdf. Accessed 06/15/04.
4. AHFS Drug Information - 2003; p. 722-724
5. USP DI - 2003; p. 2787
6. AHFS Drug Information - 2003; p. 725
7. AHFS Drug Information - 2003; p. 733-734
8. USP DI - 2003; p. 2789
9. USP DI - 2003; p. 2789
10. USP DI - 2003; p. 2788
11. USP DI - 2003; p. 2788
12. USP DI - 2003; p. 2789
13. USP DI - 2003; p. 2788
14. AHFS Drug Information - 2003; p. 735
15. USP DI - 2003; p. 2789
16. USP DI - 2003; p. 2789
17. USP DI - 2003; p. 2788
18. AHFS Drug Information - 2003; p. 735
19. AHFS Drug Information - 2003; p. 735
20. AHFS Drug Information - 2003; p. 735
21. GlaxoSmithKline - Retrovir Prescribing Information, April 2003, p. 1. Available at: http://us.gsk.com/products/assets/us_retrovir.pdf. Accessed 06/15/04.
22. USP DI - 2003; pp. 2790-2791
23. GlaxoSmithKline - Retrovir Prescribing Information, April 2003, p. 11. Available at: http://us.gsk.com/products/assets/us_retrovir.pdf. Accessed 06/15/04.
24. GlaxoSmithKline - Retrovir Prescribing Information, April 2003, p. 15. Available at: http://us.gsk.com/products/assets/us_retrovir.pdf. Accessed 06/15/04.
25. USP DI - 2003; pp. 2789-2790
26. USP DI - 2003; pp. 2789-2790
27. USP DI - 2003; pp. 2789-2790
28. USP DI - 2003; pp. 2789-2790
29. GlaxoSmithKline - Retrovir Prescribing Information, April 2003; p. 13. Available from: http://us.gsk.com/products/assets/us_retrovir.pdf. Accessed 06/15/04.
30. USP DI - 2003; pp. 2789-2790

Zidovudine



31. GlaxoSmithKline - Retrovir Prescribing Information, April 2003; p. 13. Available from: http://us.gsk.com/products/assets/us_retrovir.pdf. Accessed 06/15/04.
32. GlaxoSmithKline - Retrovir Prescribing Information, April 2003; p. 8. Available from: http://us.gsk.com/products/assets/us_retrovir.pdf. Accessed 06/15/04.
33. GlaxoSmithKline - Retrovir Prescribing Information, April 2003; p. 12. Available from: http://us.gsk.com/products/assets/us_retrovir.pdf. Accessed 06/15/04.
34. GlaxoSmithKline - Retrovir Prescribing Information, April 2003; p. 10. Available from: http://us.gsk.com/products/assets/us_retrovir.pdf. Accessed 06/15/04.
35. GlaxoSmithKline - Retrovir Prescribing Information, April 2003, p. 10. Available at: http://us.gsk.com/products/assets/us_retrovir.pdf. Accessed 06/15/04.
36. AHFS Drug Information - 2003; p. 737
37. GlaxoSmithKline - Retrovir Prescribing Information, April 2003; p. 20-21. Available from: http://us.gsk.com/products/assets/us_retrovir.pdf. Accessed 06/15/04.
38. AHFS Drug Information - 2003; p. 737
39. GlaxoSmithKline - Retrovir IV Infusion Prescribing Information, April 2003, p. 15. Available at: http://us.gsk.com/products/assets/us_retrovir_injection.pdf. Accessed 06/15/04.
40. GlaxoSmithKline - Retrovir Prescribing Information, April 2003; p. 19-20. Available from: http://us.gsk.com/products/assets/us_retrovir.pdf. Accessed 06/15/04.
41. AHFS Drug Information - 2003; p. 737
42. GlaxoSmithKline - Retrovir Prescribing Information, April 2003, p. 21-22. Available at: http://us.gsk.com/products/assets/us_retrovir.pdf. Accessed 06/15/04.
43. AHFS Drug Information - 2003; p. 737
44. GlaxoSmithKline - Retrovir Prescribing Information, April 2003, p. 21-22. Available at: http://us.gsk.com/products/assets/us_retrovir.pdf. Accessed 06/15/04.
45. AHFS Drug Information - 2003; p. 737
46. GlaxoSmithKline - Retrovir IV Infusion Prescribing Information, April 2003, p. 15. Available at: http://us.gsk.com/products/assets/us_retrovir_injection.pdf. Accessed 06/15/04.
47. ChemIDplus. - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>. Accessed 06/15/04.
48. ChemIDplus. - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>. Accessed 06/15/04.
49. ChemIDplus. - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>. Accessed 06/15/04.
50. Merck Index - 2001; p. 1809
51. Merck Index - 2001; p. 1809
52. Merck Index - 2001; p. 1809
53. GlaxoSmithKline - Retrovir Prescribing Information, April 2003, p. 2. Available at: http://us.gsk.com/products/assets/us_retrovir.pdf. Accessed 06/15/04.
54. GlaxoSmithKline - Retrovir IV Infusion Prescribing Information, April 2003, p. 15. Available at: http://us.gsk.com/products/assets/us_retrovir_injection.pdf. Accessed 06/15/04.
55. GlaxoSmithKline - Retrovir Prescribing Information, April 2003, p. 2. Available at: http://us.gsk.com/products/assets/us_retrovir.pdf. Accessed 06/15/04.
56. ChemIDplus. - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>. Accessed 06/15/04.
57. ChemIDplus. - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>. Accessed 06/15/04.
58. ChemIDplus. - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>. Accessed 06/15/04.
59. ChemIDplus. - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>. Accessed 06/15/04.
60. ChemIDplus. - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>. Accessed 06/15/04.