



**Recommendations for the Use of Antiretroviral Drugs in
Pregnant Women with HIV Infection and Interventions to Reduce
Perinatal HIV Transmission in the United States**

Downloaded from <https://aidsinfo.nih.gov/guidelines> on 5/27/2018

Visit the *AIDSinfo* website to access the most up-to-date guideline.

Register for e-mail notification of guideline updates at <https://aidsinfo.nih.gov/e-news>.

Panel's Recommendations

- Scheduled cesarean delivery at 38 weeks' gestation to minimize perinatal transmission of HIV is recommended for women with HIV RNA levels >1,000 copies/mL or unknown HIV levels near the time of delivery, irrespective of administration of antepartum antiretroviral therapy (ART) **(AII)**.
- Scheduled cesarean delivery performed solely for prevention of perinatal transmission in women receiving ART with HIV RNA ≤1,000 copies/mL is not routinely recommended due to the low rate of perinatal transmission in this group **(AII)**.
- In women with HIV RNA levels ≤1000 copies/mL, if scheduled cesarean delivery or induction is indicated, it should be performed at the standard time for obstetrical indications **(AII)**.
- In women with an HIV RNA >1,000 copies/mL or unknown HIV RNA level who present in spontaneous labor or with ruptured membranes, there is insufficient evidence to determine whether cesarean reduces the risk of perinatal HIV transmission. Management of women originally scheduled for cesarean delivery because of HIV infection who present in labor must be individualized at the time of presentation **(BII)**. In these circumstances, consultation with an expert in perinatal HIV (e.g., telephone consultation with the National Perinatal HIV/AIDS Clinical Consultation Center at (888) 448-8765) may be helpful in rapidly developing an individualized delivery plan.
- In women on ART with HIV RNA ≤1,000 copies/mL, duration of ruptured membranes is not associated with an increased risk of perinatal transmission, and vaginal delivery is recommended **(BII)**.

Rating of Recommendations: A = Strong; B = Moderate; C = Optional

Rating of Evidence: I = One or more randomized trials with clinical outcomes and/or validated laboratory endpoints; II = One or more well-designed, nonrandomized trials or observational cohort studies with long-term clinical outcomes; III = Expert opinion

Basis for Current Recommendations

Scheduled cesarean delivery, defined as cesarean delivery performed before the onset of labor and before rupture of membranes, is recommended for prevention of perinatal transmission of HIV in women with HIV RNA levels >1,000 copies/mL near delivery and for women with unknown HIV RNA levels.

This recommendation is based on findings from a multicenter, randomized clinical trial¹ and from a large individual patient data meta-analysis.² These two studies were conducted at a time when the majority of women with HIV received no antiretroviral (ARV) drugs or zidovudine as a single drug and before the availability of viral load information. Study results have since been extrapolated to make current recommendations about the mode of delivery in an era when antiretroviral therapy (ART) during pregnancy is recommended and viral load information is readily available.

HIV RNA Level of >1000 copies/mL as a Threshold for Recommendation of Scheduled Cesarean Delivery

The American Congress of Obstetricians and Gynecologists (ACOG) recommends that women with HIV RNA >1,000 copies/mL be counseled regarding the potential benefits of scheduled cesarean delivery.³ Initially, the threshold of 1,000 copies/mL was based largely on data from the Women and Infants Transmission Study, a large prospective cohort study that reported no HIV transmission among 57 women with HIV RNA levels <1,000 copies/mL.⁴ Studies reported since then have demonstrated that HIV transmission can occur in infants born to women with low viral loads.

In an analysis of 957 women with plasma viral loads ≤1,000 copies/mL, cesarean delivery (scheduled or urgent) reduced the risk of HIV transmission when adjusting for potential confounders including receipt of maternal ARV medications (AOR 0.30; *P* = 0.022); however, zidovudine alone was the regimen primarily used as prophylaxis.⁵ Among infants born to 834 women with HIV RNA ≤1,000 copies/mL receiving ARV medications, 8 (1%) were born with HIV. In a report from a comprehensive national surveillance system in the United Kingdom and Ireland, 3 (0.1%) of 2,309 and 12 (1.2%) of 1,023 infants born to women with HIV RNA levels <50 copies/mL and 50 to 999 copies/mL, respectively, were born with HIV, some of which appear to represent *in utero* transmission.⁶

Some studies demonstrate that transmission can occur even at very low HIV RNA levels. However, given the

low rate of transmission in this group, it is unclear whether scheduled cesarean delivery confers any additional benefit in reducing transmission. Furthermore, there is evidence that complication rates for cesarean deliveries are higher in women with HIV compared with women without HIV.⁷ Therefore, decisions about mode of delivery for women receiving ART with HIV RNA levels $\leq 1,000$ copies/mL should be individualized based on discussion between an obstetrician and a pregnant woman. Women should be informed that there is no evidence of benefit for scheduled cesarean delivery performed solely for prevention of perinatal transmission in women receiving ART with HIV RNA $\leq 1,000$ copies/mL and that it is not routinely recommended in this group.

Scheduled Cesarean Delivery in the Antiretroviral Therapy Era

In surveillance data from the United Kingdom and Ireland, pregnant women receiving ART (i.e., at least 3 drugs) had transmission rates of about 1%, unadjusted for mode of delivery.⁶ Given the low transmission rates achievable with use of maternal ART, the benefit of scheduled cesarean delivery is difficult to evaluate. Both the randomized clinical trial¹ and meta-analysis² documenting the benefits of cesarean delivery included mostly women who were receiving either no ARV drugs or zidovudine alone. However, other data partially address this issue.

In a report on births to women with HIV from the United Kingdom and Ireland between 2000 and 2011, perinatal transmission rates in women on ART with HIV RNA $< 1,000$ copies/mL with planned cesarean delivery (13/3814; 0.3%) were not significantly different than those in similar women with planned vaginal delivery (6/2238; 0.3%).⁸ Similarly, data from the French Perinatal Cohort showed no difference in transmission rates between vaginal delivery and planned cesarean delivery among women on ART with suppressed viral loads, 0.3% in both. For preterm deliveries with HIV RNA $< 1,000$ copies/mL, transmission rates were slightly higher among planned vaginal deliveries but the numbers were small and the differences were not statistically significant (1/9 [11.1%] vs. 1/17 [5.9%] for HIV RNA 400–1000 copies/mL; 1/39 [2.6%] vs. 1/56 [1.8%] for HIV RNA 50–400 copies/mL; 1/189 [0.5%] vs. 0/143 [0%] for HIV RNA < 50 copies/mL, for planned vaginal deliveries and elective cesarean deliveries, respectively).⁹ Therefore, no evidence to date suggests any benefit from scheduled cesarean delivery in women who have been receiving ART for several weeks and who have achieved virologic suppression.

When the delivery method selected is scheduled cesarean delivery and the maternal viral load is $> 1,000$ copies/mL, a 1-hour loading dose followed by a continuous intravenous (IV) zidovudine infusion for 2 hours (3 hours total) before scheduled cesarean delivery should be administered. In a study of the pharmacokinetics of IV zidovudine in 28 pregnant women, the ratio of cord blood-to-maternal-zidovudine levels increased significantly in women who received IV zidovudine for 3 to 6 hours compared with < 3 hours before delivery (1.0 vs. 0.55, respectively).¹⁰ This suggests that an interval of at least 3 hours may provide adequate time to reach equilibrium across the placenta, although the relationship between specific cord blood zidovudine levels or cord blood-to-maternal-zidovudine levels and efficacy in preventing perinatal transmission of HIV is unknown.

Because unscheduled cesarean delivery is performed for both maternal and fetal indications, when an unscheduled cesarean delivery is indicated in a woman who has a viral load $> 1,000$ copies/mL, consideration can be given to shortening the interval between initiation of IV zidovudine administration and delivery. For example, some experts recommend administering the 1-hour loading dose of IV zidovudine and not waiting to complete additional administration before proceeding with delivery.

Women Presenting Late in Pregnancy

Women with HIV who present late in pregnancy and are not receiving ARV drugs may not have HIV RNA results available before delivery. Without current therapy, HIV RNA levels are unlikely to be $\leq 1,000$ copies/mL at baseline. Even if ART was begun immediately, reduction in plasma HIV RNA to undetectable levels may take several weeks, depending on the **baseline viral load and** kinetics of viral decay for a particular drug regimen.¹¹⁻¹³ In this instance, scheduled cesarean delivery is likely to provide additional benefit in reducing the risk of perinatal transmission of HIV for women, unless viral suppression can be documented before 38 weeks' gestation. **Although some experts would recommend a cesarean delivery in a woman who has been virologically**

suppressed for a brief period of time (e.g., less than 2 weeks), many others would support a vaginal delivery in this scenario, as long as the plasma HIV RNA level was <1000 copies/mL by the day of delivery.

Timing of Scheduled Cesarean Delivery

For the general obstetric population, ACOG recommends that scheduled cesarean delivery not be performed before 39 weeks' gestation because of the risk of iatrogenic prematurity.^{14,15} However, in cases of cesarean delivery performed to prevent transmission of HIV, ACOG recommends scheduling cesarean delivery at 38 weeks' gestation in order to decrease the likelihood of onset of labor or rupture of membranes before delivery.³ In all women undergoing repeat cesarean delivery, the risk of any neonatal adverse event—including neonatal death, respiratory complications, hypoglycemia, newborn sepsis, or admission to the neonatal intensive care unit—is 15.3% at 37 weeks, 11.0% at 38 weeks, and 8.0% at 39 weeks.¹⁵ Gestational age should be determined by best obstetrical dating criteria, including last menstrual period and early ultrasound for dating purposes. Amniocentesis to document lung maturity should be avoided when possible in women with HIV and is rarely indicated before scheduled cesarean section for prevention of HIV transmission.

Among 1,194 infants born to mothers with HIV, 9 (1.6%) infants born vaginally had respiratory distress syndrome (RDS) compared with 18 (4.4%) infants born by scheduled cesarean delivery ($P < 0.001$). There was no statistically significant association between mode of delivery and infant RDS in an adjusted model that included infant gestational age and birth weight.¹⁶ Although newborn complications may be increased in planned births <39 weeks' gestation, the benefits of planned cesarean delivery at 38 weeks are generally thought to outweigh the risks if the procedure is performed for prevention of HIV transmission. When scheduled cesarean delivery is performed in women with HIV with an HIV RNA $\leq 1,000$ copies/mL for an indication other than decreasing HIV transmission, cesarean delivery should be scheduled based on ACOG guidelines for women without HIV.

Risk of Maternal Complications

Administration of perioperative antimicrobial prophylaxis is recommended for all women to decrease maternal infectious morbidity associated with cesarean delivery. Most studies have demonstrated that women with HIV have increased rates of postoperative complications, mostly infectious, compared with women without HIV and that risk of complications is related to degree of immunosuppression and the receipt of suppressive ART.¹⁷⁻²² Furthermore, a Cochrane review of six studies of women with HIV concluded that urgent cesarean delivery was associated with the highest risk of postpartum morbidity, scheduled cesarean delivery was intermediate in risk, and vaginal delivery had the lowest risk of morbidity.^{23,24} Complication rates in most studies^{1,25-29} were within the range reported in populations of women without HIV with similar risk factors and not of sufficient frequency or severity to outweigh the potential benefit of reduced perinatal HIV transmission. A recent U.S. study of nationally representative data from a large administrative database demonstrated that (even in the era of ART) infectious complications, surgical trauma, prolonged hospitalization, and in-hospital deaths remain higher among women with HIV compared to women without HIV.⁷ The rate of any complication associated with cesarean delivery was 117 per 1,000 deliveries among women with HIV compared with 67 per 1,000 deliveries among women without HIV. Therefore, women with HIV should be counseled regarding the specific risks associated with undergoing cesarean delivery in the setting of HIV infection.

In addition, caution should be exercised in proceeding with a cesarean delivery in circumstances where there is no clear evidence of benefit, especially in younger women who are likely to have additional pregnancies and perhaps multiple cesarean deliveries. Increased risk of abnormal placentation (e.g., placenta previa, placenta accrete, placenta increta, placenta percreta) and intrapartum hemorrhage are associated with increasing numbers of cesarean deliveries. These risks should be considered and discussed with the woman before proceeding with a cesarean delivery.^{30,31}

Managing Women Who Present in Early Labor or with Ruptured Membranes

Most studies have shown a similar risk of transmission for cesarean delivery performed for obstetric indications

after labor and membrane rupture as for vaginal delivery. In one study, the HIV transmission rate was similar in women undergoing emergency cesarean delivery and those delivering vaginally (1.6% vs. 1.9%, respectively).⁶ A meta-analysis of women with HIV, most of whom were receiving no ARV drugs or only zidovudine, demonstrated a 2% increased transmission risk for every additional hour of ruptured membranes.³² However, it is not clear how soon after the onset of labor or the rupture of membranes the benefit of cesarean delivery is lost.³³ A prospective study of 707 women in Ireland showed that among the 493 women on ART with HIV RNA levels <1,000 copies/mL, no cases of perinatal transmission occurred with membranes ruptured for up to 25 hours. Only a viral load of >10,000 copies/mL was an independent risk factor for perinatal transmission.³⁴ A prospective review of 2,398 women with HIV in the UK and Ireland, most of whom were virally suppressed, showed no association between duration of ruptured membranes and perinatal transmission in 2,116 term deliveries, regardless of viral load. Eighty-nine percent had HIV RNA levels <50 copies/mL; among the remaining 11%, 9% had HIV RNA levels 50–399 copies/mL, 1% 400–999 copies/mL, 0.4% 1000–9999 copies/mL, and 0.6% >10,000 copies/mL. Among mother-baby pairs with perinatal transmission and no evidence of *in utero* transmission, 2 had undetectable HIV RNA levels (<50 copies/ml), one had an HIV RNA level of 50–399 copies/ml, and 2 had levels >10,000 copies/ml. Among term deliveries, median duration of rupture of membranes was 3 hours 30 minutes; 71 (3.4%) had rupture of membranes >24 hours and 24 (1.1%) had rupture of membranes >48 hours. The authors concluded that obstetric care of women on ART at term with ruptured membranes should be “normalized.”^{35,36} Because it is not clear whether cesarean delivery after onset of labor reduces the risk of perinatal HIV transmission, management of women originally scheduled for cesarean delivery who present in labor must be individualized at the time of presentation. In these circumstances, consultation with an expert in perinatal HIV may be helpful. Because the delivery plan in the setting of labor must be made quickly, telephone consultation with a 24-hour, 7-day-a-week hotline (e.g., the National Perinatal HIV/AIDS Clinical Consultation Center (888) 448-8765) may be helpful in rapidly developing an individualized plan.

The woman’s oral ARV drug regimen should be continued, and IV zidovudine initiated (if previously planned) regardless of the mode of delivery.

When membrane rupture occurs before 37 weeks’ gestation, decisions about timing of delivery should be based on best obstetrical practices, considering risks to the infant of prematurity and of HIV transmission. Steroids should be given, if appropriate, to accelerate fetal lung maturity because no data exist to suggest that these recommendations need to be altered for women with HIV. When the decision is made to deliver, route of delivery should be according to obstetrical indications.

Operative Vaginal Delivery

In the past HIV was considered a relative contraindication to operative vaginal delivery with forceps or vacuum, but data from the era of ART had been lacking. Peters et al. reviewed 9,072 deliveries of women living with HIV in the UK between 2008 and 2016, where 80% of women had viral suppression from 2007 through 2011 and 90% from 2012 through 2014. Among the 3,023/3,663 vaginal deliveries with data as to whether forceps/vacuum were used, 249 (8.2%) involved operative delivery (5.6% forceps, 2.4% vacuum, 0.1% both, and 0.2% type not known). Among the 222 infants with known HIV status at 18 months of age, there was 1 case of HIV transmission with multiple possible causes and not enough evidence to confirm intrapartum transmission. The authors concluded that operative delivery is a safe option among women who are virally suppressed.³⁷

References

1. European Mode of Delivery Collaboration. Elective caesarean-section versus vaginal delivery in prevention of vertical HIV-1 transmission: a randomised clinical trial. *Lancet*. 1999;353(9158):1035-1039. Available at <http://www.ncbi.nlm.nih.gov/pubmed/10199349>.
2. International Perinatal HIV Group. The mode of delivery and the risk of vertical transmission of human immunodeficiency virus type 1--a meta-analysis of 15 prospective cohort studies. The International Perinatal HIV Group. *N Engl J Med*. 1999;340(13):977-987. Available at <http://www.ncbi.nlm.nih.gov/pubmed/10099139>.
3. American Congress of Obstetricians and Gynecologists. Committee Opinion number 234, May 2000. Scheduled

cesarean delivery and the prevention of vertical transmission of HIV infection. 2000. Available at <http://www.acog.org/Resources-And-Publications/Committee-Opinions/Committee-on-Obstetric-Practice/Scheduled-Cesarean-Delivery-and-the-Prevention-of-Vertical-Transmission-of-HIV-Infection>. Accessed October 30, 2017.

4. Garcia PM, Kalish LA, Pitt J, et al. Maternal levels of plasma human immunodeficiency virus type 1 RNA and the risk of perinatal transmission. Women and Infants Transmission Study Group. *N Engl J Med*. 1999;341(6):394-402. Available at <http://www.ncbi.nlm.nih.gov/pubmed/10432324>.
5. Ioannidis JP, Abrams EJ, Ammann A, et al. Perinatal transmission of human immunodeficiency virus type 1 by pregnant women with RNA virus loads <1000 copies/ml. *J Infect Dis*. 2001;183(4):539-545. Available at <http://www.ncbi.nlm.nih.gov/pubmed/11170978>.
6. Townsend CL, Cortina-Borja M, Peckham CS, de Ruiter A, Lyall H, Tookey PA. Low rates of mother-to-child transmission of HIV following effective pregnancy interventions in the United Kingdom and Ireland, 2000-2006. *AIDS*. 2008;22(8):973-981. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18453857>.
7. Kourtis AP, Ellington S, Pazol K, Flowers L, Haddad L, Jamieson DJ. Complications of cesarean deliveries among HIV-infected women in the United States. *AIDS*. 2014;28(17):2609-2618. Available at <http://www.ncbi.nlm.nih.gov/pubmed/25574961>.
8. Townsend CL, Byrne L, Cortina-Borja M, et al. Earlier initiation of ART and further decline in mother-to-child HIV transmission rates, 2000-2011. *AIDS*. 2014;28(7):1049-1057. Available at <http://www.ncbi.nlm.nih.gov/pubmed/24566097>.
9. Briand N, Jasseron C, Sibiude J, et al. Cesarean section for HIV-infected women in the combination antiretroviral therapies era, 2000-2010. *Am J Obstet Gynecol*. 2013;209(4):335 e331-335 e312. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23791563>.
10. Rodman JH, Flynn PM, Robbins B, et al. Systemic pharmacokinetics and cellular pharmacology of zidovudine in human immunodeficiency virus type 1-infected women and newborn infants. *J Infect Dis*. 1999;180(6):1844-1850. Available at <http://www.ncbi.nlm.nih.gov/pubmed/10558940>.
11. European Collaborative S, Patel D, Cortina-Borja M, Thorne C, Newell ML. Time to undetectable viral load after highly active antiretroviral therapy initiation among HIV-infected pregnant women. *Clin Infect Dis*. 2007;44(12):1647-1656. Available at <http://www.ncbi.nlm.nih.gov/pubmed/17516411>.
12. Aziz N, Sokoloff A, Kornak J, et al. Time to viral load suppression in antiretroviral-naive and -experienced HIV-infected pregnant women on highly active antiretroviral therapy: implications for pregnant women presenting late in gestation. *BJOG*. 2013;120(12):1534-1547. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23924192>.
13. Read PJ, Mandalia S, Khan P, et al. When should HAART be initiated in pregnancy to achieve an undetectable HIV viral load by delivery? *AIDS*. 2012;26(9):1095-1103. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22441248>.
14. American College of Obstetricians and Gynecologists. ACOG Practice Bulletin No. 97: Fetal lung maturity. *Obstet Gynecol*. 2008;112(3):717-726. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18757686>.
15. Tita AT, Landon MB, Spong CY, et al. Timing of elective repeat cesarean delivery at term and neonatal outcomes. *N Engl J Med*. 2009;360(2):111-120. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19129525>.
16. Livingston EG, Huo Y, Patel K, et al. Mode of delivery and infant respiratory morbidity among infants born to HIV-1-infected women. *Obstet Gynecol*. 2010;116(2 Pt 1):335-343. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20664394>.
17. Grubert TA, Reindell D, Kastner R, Lutz-Friedrich R, Belohradsky BH, Dathe O. Complications after caesarean section in HIV-1-infected women not taking antiretroviral treatment. *Lancet*. 1999;354(9190):1612-1613. Available at <http://www.ncbi.nlm.nih.gov/pubmed/10560681>.
18. Maiques-Montesinos V, Cervera-Sanchez J, Bellver-Pradas J, Abad-Carrascosa A, Serra-Serra V. Post-cesarean section morbidity in HIV-positive women. *Acta Obstet Gynecol Scand*. 1999;78(9):789-792. Available at <http://www.ncbi.nlm.nih.gov/pubmed/10535342>.
19. Rodriguez EJ, Spann C, Jamieson D, Lindsay M. Postoperative morbidity associated with cesarean delivery among human immunodeficiency virus-seropositive women. *Am J Obstet Gynecol*. 2001;184(6):1108-1111. Available at <http://www.ncbi.nlm.nih.gov/pubmed/11349171>.
20. Semprini AE, Castagna C, Ravizza M, et al. The incidence of complications after caesarean section in 156 HIV-positive

- women. *AIDS*. 1995;9(8):913-917. Available at <http://www.ncbi.nlm.nih.gov/pubmed/7576327>.
21. Urbani G, de Vries MM, Cronje HS, Niemand I, Bam RH, Beyer E. Complications associated with cesarean section in HIV-infected patients. *Int J Gynaecol Obstet*. 2001;74(1):9-15. Available at <http://www.ncbi.nlm.nih.gov/pubmed/11430935>.
 22. Vimercati A, Greco P, Loverro G, Lopalco PL, Pansini V, Selvaggi L. Maternal complications after caesarean section in HIV infected women. *Eur J Obstet Gynecol Reprod Biol*. 2000;90(1):73-76. Available at <http://www.ncbi.nlm.nih.gov/pubmed/10767514>.
 23. Read JS, Newell MK. Efficacy and safety of cesarean delivery for prevention of mother-to-child transmission of HIV-1. *Cochrane Database Syst Rev*. 2005(4):CD005479. Available at <http://www.ncbi.nlm.nih.gov/pubmed/16235405>.
 24. Livingston EG, Huo Y, Patel K, et al. Complications and route of delivery in a large cohort study of HIV-1-infected women-IMPAACT P1025. *J Acquir Immune Defic Syndr*. 2016;73(1):74-82. Available at <https://www.ncbi.nlm.nih.gov/pubmed/27082506>.
 25. Faucher P, Batallan A, Bastian H, et al. Management of pregnant women infected with HIV at Bichat Hospital between 1990 and 1998: analysis of 202 pregnancies. *Gynecol Obstet Fertil*. 2001;29(3):211-225. Available at <http://www.ncbi.nlm.nih.gov/pubmed/11300046>.
 26. Fiore S, Newell ML, Thorne C, European HIV in Obstetrics Group. Higher rates of post-partum complications in HIV-infected than in uninfected women irrespective of mode of delivery. *AIDS*. 2004;18(6):933-938. Available at <http://www.ncbi.nlm.nih.gov/pubmed/15060441>.
 27. Marcollet A, Goffinet F, Firtion G, et al. Differences in postpartum morbidity in women who are infected with the human immunodeficiency virus after elective cesarean delivery, emergency cesarean delivery, or vaginal delivery. *Am J Obstet Gynecol*. 2002;186(4):784-789. Available at <http://www.ncbi.nlm.nih.gov/pubmed/11967508>.
 28. Read JS, Tuomala R, Kpamegan E, et al. Mode of delivery and postpartum morbidity among HIV-infected women: the women and infants transmission study. *J Acquir Immune Defic Syndr*. 2001;26(3):236-245. Available at <http://www.ncbi.nlm.nih.gov/pubmed/11242196>.
 29. Watts DH, Lambert JS, Stiehm ER, et al. Complications according to mode of delivery among human immunodeficiency virus-infected women with CD4 lymphocyte counts of < or = 500/microL. *Am J Obstet Gynecol*. 2000;183(1):100-107. Available at <http://www.ncbi.nlm.nih.gov/pubmed/10920316>.
 30. Silver RM, Landon MB, Rouse DJ, et al. Maternal morbidity associated with multiple repeat cesarean deliveries. *Obstet Gynecol*. 2006;107(6):1226-1232. Available at <https://www.ncbi.nlm.nih.gov/pubmed/16738145>.
 31. Greenbaum S, Wainstock T, Dukler D, Leron E, Erez O. Underlying mechanisms of retained placenta: Evidence from a population based cohort study. *Eur J Obstet Gynecol Reprod Biol*. 2017;216:12-17. Available at <https://www.ncbi.nlm.nih.gov/pubmed/28692888>.
 32. International Perinatal HIV Group. Duration of ruptured membranes and vertical transmission of HIV-1: a meta-analysis from 15 prospective cohort studies. *AIDS*. 2001;15(3):357-368. Available at <http://www.ncbi.nlm.nih.gov/pubmed/11273216>.
 33. Jamieson DJ, Read JS, Kourtis AP, Durant TM, Lampe MA, Dominguez KL. Cesarean delivery for HIV-infected women: recommendations and controversies. *Am J Obstet Gynecol*. 2007;197(3 Suppl):S96-100. Available at <http://www.ncbi.nlm.nih.gov/pubmed/17825656>.
 34. Cotter AM, Brookfield KF, Duthely LM, Gonzalez Quintero VH, Potter JE, O'Sullivan MJ. Duration of membrane rupture and risk of perinatal transmission of HIV-1 in the era of combination antiretroviral therapy. *Am J Obstet Gynecol*. 2012;207(6):482 e481-485. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23103331>.
 35. Peters H, Byrne L, De Ruiter A, et al. Duration of ruptured membranes and mother-to-child HIV transmission: a prospective population-based surveillance study. *BJOG*. 2015. Available at <http://www.ncbi.nlm.nih.gov/pubmed/26011825>.
 36. Eppes C. Is it time to leave the avoidance of rupture of membranes for women infected with HIV and receiving cART in the past? *BJOG*. 2015. Available at <http://www.ncbi.nlm.nih.gov/pubmed/25998194>.
 37. Peters H, Francis K, Harding K, Tookey PA, Thorne C. Operative vaginal delivery and invasive procedures in pregnancy among women living with HIV. *Eur J Obstet Gynecol Reprod Biol*. 2017;210:295-299. Available at <https://www.ncbi.nlm.nih.gov/pubmed/28092853>.