



## **Guidelines for the Use of Antiretroviral Agents in Adults and Adolescents with HIV**

Downloaded from <https://aidsinfo.nih.gov/guidelines> on 9/13/2019

Visit the *AIDSinfo* website to access the most up-to-date guideline.

Register for e-mail notification of guideline updates at <https://aidsinfo.nih.gov/e-news>.

## Co-Receptor Tropism Assays (Last updated October 25, 2018; last reviewed October 25, 2018)

### Panel's Recommendations

- A co-receptor tropism assay should be performed whenever the use of a CCR5 co-receptor antagonist is being considered (AI).
- Co-receptor tropism testing is recommended for patients who exhibit virologic failure on a CCR5 antagonist (BIII).
- A phenotypic tropism assay is preferred to determine HIV-1 co-receptor usage (AI).
- A genotypic tropism assay should be considered as an alternative test to predict HIV-1 co-receptor usage (BII).
- A proviral DNA tropism assay can be utilized for patients with undetectable HIV-1 RNA when a CCR5 antagonist is considered for use in a new regimen (e.g., as part of a regimen switch or simplification) (BII).

**Rating of Recommendations:** A = Strong; B = Moderate; C = Optional

**Rating of Evidence:** I = Data from randomized controlled trials; II = Data from well-designed nonrandomized trials or observational cohort studies with long-term clinical outcomes; III = Expert opinion

HIV enters cells by a complex process that involves sequential attachment to the CD4 T lymphocyte (CD4) receptor followed by binding to either the CCR5 or CXCR4 molecules and fusion of the viral and cellular membranes.<sup>1</sup> CCR5 co-receptor antagonists prevent HIV entry into target cells by binding to the CCR5 receptors.<sup>2</sup> Phenotypic and genotypic assays have been developed that can determine or predict the co-receptor tropism (i.e., use of CCR5, CXCR4, or both as either dual-tropic virus or a mixed population of viruses referred to for purposes of assay results as dual/mixed [D/M]) of the patient's dominant virus population. An older generation assay (Trofile,<sup>®</sup> Monogram Biosciences, Inc., South San Francisco, CA) was used to screen patients who were participating in clinical trials that led to the approval of maraviroc (MVC), the only CCR5 antagonist currently available. The assay has been improved and is now available with enhanced sensitivity. In addition, genotypic assays to predict co-receptor usage are commercially available.

During acute/recent infection, the vast majority of patients harbor a CCR5-utilizing virus (R5 virus), which suggests that the R5 variant is preferentially transmitted; however, up to 19% of individuals with acute/recent infection can harbor CXCR4-tropic virus.<sup>3-5</sup> Viruses in many untreated patients eventually exhibit a shift in co-receptor tropism from CCR5 usage to either CXCR4 usage or D/M tropism. This shift is temporally associated with a more rapid decline in CD4 counts,<sup>6,7</sup> but whether this tropism shift is a cause or a consequence of progressive immunodeficiency remains undetermined.<sup>1</sup> Antiretroviral-treated patients with extensive drug resistance or persistently high-level viremia are more likely to harbor CXCR4- or D/M-tropic variants than untreated patients with comparable CD4 counts.<sup>8,9</sup> The prevalence of CXCR4- or D/M-tropic variants increases to more than 50% in treated patients who have CD4 counts <100 cells/mm<sup>3</sup>.<sup>8,10</sup> Since CXCR4-tropic viruses may be present at initial presentation or a patient may shift to CXCR4-tropism over the course of infection, co-receptor tropism should always be assessed prior to the use of CCR5 antagonists for treatment. Once a patient has ever been documented with detectable CXCR4- or D/M-tropic virus, it is assumed that such viruses will always be present. CCR5 co-receptor antagonists will no longer be active for that patient and should not be used.

### Phenotypic Assays

Phenotypic assays characterize the co-receptor usage of plasma-derived virus. These assays involve the generation of laboratory viruses that express patient-derived envelope proteins (i.e., gp120 and gp41). These pseudoviruses, which are replication-defective, are used to infect target cell lines that express either CCR5 or CXCR4.<sup>11,12</sup> Using the Trofile<sup>®</sup> assay, the co-receptor tropism of the patient-derived virus is confirmed by testing the susceptibility of the virus to specific CCR5 or CXCR4 inhibitors *in vitro*. This assay takes about 2 weeks to perform and requires a plasma HIV RNA level  $\geq 1,000$  copies/mL.

The performance characteristics of these assays have evolved. Most, if not all, patients enrolled in premarketing clinical trials of MVC and other CCR5 antagonists were screened with an earlier, less sensitive version of the Trofile<sup>®</sup> assay.<sup>12</sup> This earlier assay failed to routinely detect low levels of CXCR4-utilizing variants. As a consequence, some patients enrolled in these clinical trials harbored low levels of such variants at baseline, which were below the assay limit of detection, and these patients exhibited rapid virologic failure after initiation of a CCR5 antagonist.<sup>13</sup> The assay has been improved and is now able to detect lower levels of CXCR4-utilizing viruses. *In vitro*, the assay can detect CXCR4-utilizing clones with 100% sensitivity when those clones represent 0.3% or more of the virus population.<sup>14</sup> Although this more sensitive assay has had limited use in prospective clinical trials, it is now the only assay that is commercially available. For unclear reasons, a minority of samples cannot be successfully phenotyped with either generation of the Trofile<sup>®</sup> assay.

In patients with an undetectable viral load or **detectable plasma HIV RNA <1,000 copies/mL**, phenotypic co-receptor usage can be determined using proviral DNA obtained from peripheral blood mononuclear cells (e.g., Trofile<sup>®</sup> DNA, Monogram Sciences); however, the clinical utility of this assay remains to be determined.<sup>15</sup>

### ***Genotypic Assays***

Genotypic determination of HIV-1 co-receptor usage is based on sequencing of the V3-coding region of HIV-1 *env*, the principal determinant of co-receptor usage. A variety of algorithms and bioinformatics programs can be used to predict co-receptor usage from the V3 sequence.<sup>16</sup> When compared to the phenotypic assay, genotypic methods show high specificity (~90%) but only modest sensitivity (~50% to 75%) for the presence of a CXCR4-utilizing virus. Studies in which V3 genotyping was performed on samples from patients screened for clinical trials of MVC suggest that genotyping performed as well as phenotyping in predicting the response to MVC.<sup>17-19</sup> An important caveat is that the majority of patients who received MVC were first shown to have R5 virus by a phenotypic assay (Trofile<sup>®</sup>). Consequently, the opportunity to assess treatment response to MVC in patients whose virus was considered R5 by genotype but D/M or X4 by phenotype was limited to a relatively small number of patients. **Other studies have also demonstrated relatively high concordance between genotypic- and phenotypic-assessed tropism;<sup>20,21</sup> however, there is variability between different genotypic platforms.<sup>22</sup>**

Given these performance characteristics, genotypic tropism assays may not be sufficiently robust to completely rule out the presence of an X4 or D/M variant;<sup>23</sup> therefore, the Panel preferentially recommends phenotypic testing. **Based on accessibility, capacity, logistics, and cost, European guidelines currently include genotypic testing as an equivalent option to phenotypic testing when determining co-receptor usage among patients with HIV RNA >1,000 copies/mL and preferentially for those with HIV RNA ≤1,000 copies/mL.<sup>24</sup>**

**HIV-1 proviral DNA genotypic tropism testing is available for patients with HIV RNA <1,000 copies/mL. These assays evaluate the HIV-1 proviral DNA integrated within infected cells for CXCR4-utilizing viral strains.<sup>25</sup> As discussed above, caution is advised when using such assays, as their detection limit, concordance with plasma HIV RNA tropism, and clinical utility are not yet fully determined.**

### ***Use of Assays to Determine Co-receptor Usage in Clinical Practice***

An assay for HIV-1 co-receptor usage should be performed whenever the use of a CCR5 antagonist is being considered (**AI**). This is true even in the setting of prior tropism testing showing CCR5 usage, as viral evolution may occur over the course of infection. In addition, because virologic failure may occur due to a shift from CCR5-using to CXCR4-using virus, testing for co-receptor usage is recommended in patients who exhibit virologic failure on a CCR5 antagonist (**BIII**). Virologic failure may also be caused by resistance of a CCR5-using virus to a CCR5 antagonist, but such resistance is uncommon. Compared to genotypic testing, phenotypic testing has more evidence supporting its utility. Therefore, a phenotypic test for co-receptor usage is generally preferred (**AI**). However, because phenotypic testing is more expensive, requires more time to

perform, and may have logistic challenges, a genotypic test to predict HIV-1 co-receptor usage should be considered as an alternative test **(BII)**.

As with HIV resistance testing, the results of all prior tropism tests should be obtained. If CXCR4-utilizing or D/M-tropic viruses have ever been detected previously, then repeat testing is not necessary and a CCR5 co-receptor antagonist **should not be used**.

If a CCR5 co-receptor antagonist is being considered in a patient with an undetectable HIV RNA (e.g., in cases of regimen simplification or a toxicity-related switch), a proviral DNA tropism assay can be utilized **(BII)**.<sup>26-28</sup> If CXCR4-utilizing or D/M-tropic viruses are detected, then the CCR5 co-receptor antagonist **should not be used**.

## References

1. Moore JP, Kitchen SG, Pugach P, Zack JA. The CCR5 and CXCR4 coreceptors--central to understanding the transmission and pathogenesis of human immunodeficiency virus type 1 infection. *AIDS Res Hum Retroviruses*. 2004;20(1):111-126. Available at: [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=15000703](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15000703).
2. Fatkenheuer G, Pozniak AL, Johnson MA, et al. Efficacy of short-term monotherapy with maraviroc, a new CCR5 antagonist, in patients infected with HIV-1. *Nat Med*. 2005;11(11):1170-1172. Available at: [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=16205738](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16205738).
3. Zhu T, Mo H, Wang N, et al. Genotypic and phenotypic characterization of HIV-1 patients with primary infection. *Science*. 1993;261(5125):1179-1181. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8356453>.
4. Brumme ZL, Goodrich J, Mayer HB, et al. Molecular and clinical epidemiology of CXCR4-using HIV-1 in a large population of antiretroviral-naïve individuals. *J Infect Dis*. 2005;192(3):466-474. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15995960>.
5. Raymond S, Nicot F, Saune K, et al. Brief report: HIV-1 tropism during primary Infections in France: 1996-2014. *J Acquir Immune Defic Syndr*. 2016;72(4):376-379. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26959188>.
6. Connor RI, Sheridan KE, Ceradini D, Choe S, Landau NR. Change in coreceptor use correlates with disease progression in HIV-1-infected individuals. *J Exp Med*. 1997;185(4):621-628. Available at: [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=9034141](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=9034141).
7. Koot M, Keet IP, Vos AH, et al. Prognostic value of HIV-1 syncytium-inducing phenotype for rate of CD4+ cell depletion and progression to AIDS. *Ann Intern Med*. 1993;118(9):681-688. Available at: [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=8096374](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=8096374).
8. Hunt PW, Harrigan PR, Huang W, et al. Prevalence of CXCR4 tropism among antiretroviral-treated HIV-1-infected patients with detectable viremia. *J Infect Dis*. 2006;194(7):926-930. Available at: [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=16960780](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16960780).
9. Agwu AL, Yao TJ, Eshleman SH, et al. Phenotypic coreceptor tropism in perinatally HIV-infected youth failing antiretroviral therapy. *Pediatr Infect Dis J*. 2016;35(7):777-781. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27078121>.
10. Wilkin TJ, Su Z, Kuritzkes DR, et al. HIV type 1 chemokine coreceptor use among antiretroviral-experienced patients screened for a clinical trial of a CCR5 inhibitor: AIDS Clinical Trial Group A5211. *Clin Infect Dis*. 2007;44(4):591-595. Available at: [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=17243065](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17243065).
11. Trouplin V, Salvatori F, Cappello F, et al. Determination of coreceptor usage of human immunodeficiency virus type 1 from patient plasma samples by using a recombinant phenotypic assay. *J Virol*. 2001;75(1):251-259. Available at: [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=11119595](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11119595).
12. Whitcomb JM, Huang W, Fransen S, et al. Development and characterization of a novel single-cycle recombinant-virus assay to determine human immunodeficiency virus type 1 coreceptor tropism. *Antimicrob Agents Chemother*. 2007;51(2):566-575. Available at: <http://www.ncbi.nlm.nih.gov/entrez/query>.

[fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=17116663](https://www.ncbi.nlm.nih.gov/pubmed/17116663).

13. Westby M, Lewis M, Whitcomb J, et al. Emergence of CXCR4-using human immunodeficiency virus type 1 (HIV-1) variants in a minority of HIV-1-infected patients following treatment with the CCR5 antagonist maraviroc is from a pretreatment CXCR4-using virus reservoir. *J Virol*. 2006;80(10):4909-4920. Available at: [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=16641282](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16641282).
14. Trinh L, Han D, Huang W, et al. Technical validation of an enhanced sensitivity Trofile HIV coreceptor tropism assay for selecting patients for therapy with entry inhibitors targeting CCR5. *Antivir Ther*. 2008;13(Suppl 3):A128.
15. Toma J, Frantzell A, Cook J, et al. Phenotypic determination of HIV-1 coreceptor tropism using cell-associated DNA derived from blood samples. Presented at: Conference on Retroviruses and Opportunistic Infections. 2010. San Francisco, CA.
16. Garrido C, Roulet V, Chueca N, et al. Evaluation of eight different bioinformatics tools to predict viral tropism in different human immunodeficiency virus type 1 subtypes. *J Clin Microbiol*. 2008;46(3):887-891. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18199789>.
17. McGovern RA, Thielen A, Mo T, et al. Population-based V3 genotypic tropism assay: a retrospective analysis using screening samples from the A4001029 and MOTIVATE studies. *AIDS*. 2010;24(16):2517-2525. Available at: [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=20736814](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20736814).
18. McGovern RA, Thielen A, Portsmouth S, et al. Population-based sequencing of the V3-loop can predict the virological response to maraviroc in treatment-naïve patients of the MERIT trial. *J Acquir Immune Defic Syndr*. 2012;61(3):279-286. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23095934>.
19. Archer J, Weber J, Henry K, et al. Use of four next-generation sequencing platforms to determine HIV-1 coreceptor tropism. *PLoS One*. 2012;7(11):e49602. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23166726>.
20. Heger E, Kaiser R, Knops E, et al. Results of the first international HIV-1 coreceptor proficiency panel test. *J Clin Virol*. 2017;93:53-56. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28633097>.
21. Kagan RM, Johnson EP, Siaw MF, et al. Comparison of genotypic and phenotypic HIV type 1 tropism assay: results from the screening samples of Cenicriviroc Study 202, a randomized phase II trial in treatment-naïve subjects. *AIDS Res Hum Retroviruses*. 2014;30(2):151-159. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23875707>.
22. Swenson LC, Dong WW, Mo T, et al. Use of cellular HIV DNA to predict virologic response to maraviroc: performance of population-based and deep sequencing. *Clin Infect Dis*. 2013;56(11):1659-1666. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23429552>.
23. Lin NH, Kuritzkes DR. Tropism testing in the clinical management of HIV-1 infection. *Curr Opin HIV AIDS*. 2009;4(6):481-487. Available at: [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=20048714](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20048714).
24. Vandekerckhove LP, Wensing AM, Kaiser R, et al. European guidelines on the clinical management of HIV-1 tropism testing. *Lancet Infect Dis*. 2011;11(5):394-407. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21429803>.
25. Fabeni L, Berno G, Svicher V, et al. Genotypic tropism testing in HIV-1 proviral DNA can provide useful information at low-level viremia. *J Clin Microbiol*. 2015;53(9):2935-2941. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26135872>.
26. Vitiello P, Brudney D, MacCartney M, et al. Responses to switching to maraviroc-based antiretroviral therapy in treated patients with suppressed plasma HIV-1-RNA load. *Intervirology*. 2012;55(2):172-178. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22286889>.
27. Bonjoch A, Pou C, Perez-Alvarez N, et al. Switching the third drug of antiretroviral therapy to maraviroc in aviraemic subjects: a pilot, prospective, randomized clinical trial. *J Antimicrob Chemother*. 2013;68(6):1382-1387. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23354282>.
28. Pett SL, Amin J, Horban A, et al. Maraviroc, as a switch option, in HIV-1-infected individuals with stable, well-controlled HIV replication and R5-tropic virus on their first nucleoside/nucleotide reverse transcriptase inhibitor plus ritonavir-boosted protease inhibitor regimen: Week 48 results of the randomized, multicenter MARCH Study. *Clin Infect Dis*. 2016;63(1):122-132. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27048747>.